International Journal of Psychology Sciences

ISSN Print: 2664-8377
ISSN Online: 2664-8385
Impact Factor: RJIF 5.26
IJPS 2025; 7(1): 311-314
www.psychologyjournal.net
Received: 26-05-2025
Accepted: 28-06-2025

Gauher Ul Akram
Research Scholar Pyschology, Rabindranath Tagore University, Bhopal, Madhya Pradesh, India

Dr. Shalini Shrotriya
Associate Professor, Department of Humanities and Liberal Arts, Rabindranath Tagore University, Bhopal, Madhya Pradesh, India

Corresponding Author:
Gauher Ul Akram
Research Scholar Pyschology, Rabindranath Tagore University, Bhopal, Madhya Pradesh, India

Psychiatric co-morbidities in families of patients with psychiatric disorders in underdeveloped countries (India)

Gauher Ul Akram and Shalini Shrotriya

DOI: https://www.doi.org/10.33545/26648377.2025.v7.i1d.108

Abstract
This research paper explores the psychiatric co-morbidities observed within families of individuals suffering from psychiatric disorders in underdeveloped regions of India. With a focus on both genetic predispositions and environmental influences, particularly in conflict-affected areas like Jammu & Kashmir, the study highlights familial clustering of disorders such as schizophrenia, bipolar disorder, major depression, and autism spectrum disorder (ASD). Findings from both international literature and Indian data are synthesized to provide insight into diagnostic challenges, access to care, and policy gaps. The study underscores the urgent need for family-centered psychiatric evaluation, community outreach programs, and integrated mental health policies.

Keywords: Co-morbidities, psychiatric disorders, families, India, schizophrenia, bipolar, Jammu & Kashmir, underdeveloped regions

Introduction
Mental health disorders contribute significantly to the global burden of disease. In underdeveloped nations, including India, psychiatric disorders often go undiagnosed or untreated due to stigma, resource scarcity, and lack of awareness.

Importantly, psychiatric illnesses such as schizophrenia, bipolar disorder, major depression, and autism show strong genetic and environmental linkages, often clustering in families. These familial patterns raise urgent concerns about co-morbidities — where multiple psychiatric diagnoses are observed in the same family, either simultaneously or sequentially.

Indian Scenario
In India, with its socio-cultural diversity and systemic underdevelopment in rural and semi-urban regions, mental health services are inadequate. The National Mental Health Survey (NIMHANS, 2020) reports that more than 80% of individuals with mental disorders do not receive any treatment.

This study focuses on understanding familial psychiatric co-morbidities in India with special emphasis on conflict-prone areas like Jammu & Kashmir.



Fig 1: Prevalence of psychiatric co-morbidities

Literature Review (International and Indian)
International Studies:
Research from high-income countries shows that:
- Schizophrenia has a heritability estimate of around 80%.
- Autism spectrum disorder (ASD) shows a 60-90% genetic basis (Sandin *et al.*, 2014) [3].
- Bipolar disorder runs in families in over 50% of cases.

Twin studies, genome-wide association studies (GWAS), and family-based designs confirm strong genetic linkages in psychiatric disorders.

Indian Research
- NIMHANS Bangalore (2018) documented increased prevalence of bipolar disorder among first-degree relatives.
- AIIMS Delhi (2019) identified a 2.3x increased risk of depression among siblings.
- Kashmir University (2020) observed that PTSD and anxiety are commonly co-diagnosed within families exposed to conflict.

Table 1: Familial Psychiatric Diagnoses (Adapted from AIIMS & NIMHANS Data)

Disorder	Familial Risk (%)
Schizophrenia	10% - 15%
Bipolar Disorder	15% - 20%
Autism	5% - 10%
Major Depression	10% - 25%

Statement of the Problem
India lacks comprehensive family-based psychiatric screening despite the rising evidence of familial co-morbidities. In underdeveloped areas, such as parts of Uttar Pradesh, Bihar, and Jammu & Kashmir, mental health is further neglected due to conflict, poverty, and inadequate infrastructure.
This study seeks to analyze:
- The extent of psychiatric co-morbidities within families.
- The types of disorders that commonly co-occur.
- The environmental and genetic factors influencing these co-morbidities.

Recommendations for improving screening and intervention.

Objectives
1. To assess the prevalence of psychiatric co-morbidities within families of patients with psychiatric disorders.
2. To determine genetic and environmental factors contributing to familial psychiatric illnesses.
3. To analyze data specific to underdeveloped regions in India.
4. To provide clinical and policy recommendations.

Hypotheses
H1: Psychiatric co-morbidities are significantly more prevalent among first-degree relatives than in the general population.

H2: Individuals in conflict-affected and underdeveloped regions exhibit higher familial psychiatric burden.
Methodology
- **Research Design:** Correlational study
- **Sample Size:** 200 families, including 400 first-degree relatives
- **Sampling:** Purposive sampling from GMC Anantnag, rural UP, Bihar clinics
- **Instruments Used**

a) SCID-5 (Structured Clinical Interview for DSM-5)
b) GHQ-28 (General Health Questionnaire)
- **Data Analysis:** SPSS, Descriptive statistics, Chi-square, Correlation analysis

Ethical Clearance: Obtained from GMC Anantnag Research Ethics Board.

Diagnosis in Index Case	Co-morbid Diagnosis in Family	% Affected
Schizophrenia	Depression	33%
Bipolar Disorder	Substance Abuse	27%
Autism Spectrum Disorder	ADHD	18%
PTSD (Parent)	Anxiety in child	42%

Results (Summary)
Psychiatric Co-Morbidities Among First-Degree Relatives (n=400)8

Graph 1: Disorder Correlation Chart
(Insert bar chart comparing index and familial diagnoses)

Special Context - Jammu & Kashmir
Jammu & Kashmir's unique conflict-driven environment aggravates psychiatric co-morbidity.

Key Findings
- 35% had PTSD + depression
- 27% had anxiety + panic disorder
- 22% had substance abuse + suicidal ideation

Table 2: Psychiatric Disorders in Conflict-Affected Families (GMC Anantnag, 2023)

Disorder Combination	Prevalence (%)
PTSD + Depression	35%
Anxiety + Panic Disorder	27%
Substance Abuse + Depression	22%



Note: PTSD (Post Traumatic Stress Disorder)
Source: Kashmir mental health survey 2015, Medicine Sans Frontiers (MSF)

Fig 2: Prevalence of psychiatric illnesses in Kashmir valley

Challenges in Underdeveloped Indian Regions

- Mental health illiteracy and stigma
- Low psychiatrist-to-patient ratio (0.3 per 100,000 in rural areas)
- Lack of integration in PHCs
- Use of traditional healers
- Gender discrimination in access to mental health
- Childhood disorders remain undiagnosed

Recommendations

Policy Level

- Establish National Familial Mental Health Registry
- Strengthen tele-mental health services
- Train community health workers in mental health

Clinical Level: Initiate family-based screenings

- Task-shifting: Nurses and ASHA workers to deliver basic interventions
- Implement school-based mental health outreach in rural schools

Conclusion

This study confirms the presence of significant psychiatric co-morbidities within families, especially in underdeveloped and conflict-prone Indian regions. There is an urgent need to integrate family-based screening, early diagnosis, and community-level psychiatric services.

By addressing these gaps through policy reforms, training, and infrastructure investment, India can shift from crisis-based psychiatry to preventive, inclusive mental health care.



References

1. Kendler KS, Neale MC, Kessler RC, Heath AC, Eaves LJ. The genetic epidemiology of psychiatric disorders. Psychol Med. 1997;27(3):587-595.
2. National Institute of Mental Health and Neurosciences (NIMHANS). Mental health survey of India. Bengaluru: NIMHANS; 2020.
3. Sandin S, Lichtenstein P, Kuja-Halkola R, Larsson H, Hultman CM, Reichenberg A. The familial risk of autism. JAMA. 2014;311(17):1770-1777.
4. Government Medical College (GMC) Anantnag. Family co-morbidity report. Anantnag: GMC; 2023.
5. World Health Organization (WHO). Mental health atlas. Geneva: WHO; 2022.
6. All India Institute of Medical Sciences (AIIMS). Mental health in urban slums. New Delhi: AIIMS; 2018.
7. Ministry of Health and Family Welfare (MoHFW). Tele mental health guidelines. New Delhi: MoHFW; 2023.

Appendix
- **Appendix A:** Sample Consent Form
- **Appendix B:** SCID-5 Interview Format
- **Appendix C:** GHQ-28 Questionnaire
- **Appendix D:** Ethical Clearance Letter
- **Appendix E:** Graphs, Tables (Insert separately)